



## Response of Soil Application of Zinc Sulphate and Ferrous Sulphate on Leaf Nutrient Status of Guava Plants cv. Hisar Safeda under High Density Planting

Rupakshi\* and Satpal Baloda

Department of Horticulture, CCSHAU, Hisar, Haryana

\*Email : [dr.rupakshi01@gmail.com](mailto:dr.rupakshi01@gmail.com)

### Abstract

The aim of present work was to evaluate leaf nutrient status of 8 years old guava plants pertaining to soil application of zinc sulphate and ferrous sulphate. Uniform plants planted at 6m X 2m spacing at Research Orchard of Department of Horticulture, CCS Haryana Agricultural University, Hisar were selected for the investigation. Maximum total nitrogen and total zinc content was noted with the application of 90g zinc sulphate. While for ferrous sulphate treatments, maximum total nitrogen and total zinc content was analysed with the application of 20g ferrous sulphate. Maximum total phosphorus and total potassium content was observed under control treatment under zinc sulphate as well as ferrous sulphate treatments. Highest iron content was noted with application of 30g zinc sulphate and 30g ferrous sulphate treatments.

**Key words :** soil application, zinc sulphate, ferrous sulphate, guava, leaf nutrients.


### Introduction

Guava (*Psidium guajava* L.), a member of family Myrtaceae, native to tropical America, the apple of the tropics, is one of the most common fruits in India. It claims to be the fourth most important fruit in area and production after mango, banana and citrus. Being very hardy, it gives an assured crop even with very little care. Fertilizers applied by growers not only affect plant growth, yield or fruit quality, but also leaf nutrient composition. The analyses of total amounts of nutrients in selected leaves or leaf parts have become an important tool in evaluating fertility status of a number of plants. They have been especially useful to determine need and composition of fertilizer side dressing and foliar sprays. (1, 2). It has been noticed that guava suffers severely from deficiency of micronutrients which reduce the quality of fruits. Knowledge of the chemical mineral composition of the leaves provide information to optimize the fertilization program to maximize efficient production and maintain soil fertility. Hence, the present study was undertaken to examine the leaf nutrient status of guava plant after soil application of zinc sulphate and ferrous sulphate.

### Materials and Methods

The work was undertaken at the Research Orchard of Department of Horticulture, CCS Haryana Agricultural University, Hisar to study the effect of different levels of zinc sulphate (zero, 30, 60 and 90g/plant) and ferrous sulphate (zero, 10, 20 and 30g/plant) on leaf nutrient status of guava cv. Hisar Safeda. Uniform guava plants of 8 years age planted at 6m X 2m spacing were selected for this study. Leaf samples were collected during August

month from middle of non-fruiting branches. Forty to fifty leaves were taken from each tree. These were grinded using grinder and the powder thus formed was stored in clean polythene bags. These powdered leaf samples were used for digestion.



Element	Method of Estimation
Total Nitrogen (%)	Colorimetric or Nessler's method proposed by (3)
Total Phosphorus (%)	Vanado-molybdophosphoric yellow color method proposed by (4)
Total Potassium (%)	Flame photometry
Total micronutrients [Zn and Fe (ppm)]	Atomic Absorption Spectrophotometer (AAS)

## Results and Discussion

**Nitrogen content in leaf :** Persual from the data shown in Table-1 reveals that nitrogen content in guava leaf improved significantly on application of zinc sulphate. Maximum nitrogen content (1.74%) was observed from the leaves of plants receiving 90 g zinc sulphate treatment, while minimum (1.68%) was observed with control treatment. It may be due to the fact that increase in zinc sulphate leads to reducing nitrogen concentration in leaves. Similar results were obtained by (5) in apple, where application of zinc (soil and foliar) significantly increased nitrogen concentration of leaf. In contradiction to the present results, non-significant results were obtained by (6) in apple and (7) in peach. They reported non-significant effect of zinc sulphate on leaf nitrogen composition. Nitrogen content in leaves was significantly affected with the application of ferrous sulphate. Maximum nitrogen concentration (1.74%) was obtained from application of 20 g ferrous sulphate, which was at par with 30 g ferrous sulphate and significantly higher than all other treatments, while minimum (1.69%) was noted under control treatment. In opposite to it, (8) revealed that nitrogen concentration remain unaffected with iron application under greenhouse as well as field conditions. No significant effect on nitrogen content was found due to interaction between zinc sulphate and ferrous sulphate.

**Phosphorus content in leaf :** The phosphorus content in guava leaves was significantly affected due to application of zinc sulphate (Table-1). Maximum phosphorus content (0.18%) was observed under control treatment which was at par with 30 g zinc sulphate as well as 90 g zinc sulphate and minimum (0.16%) from plant receiving 60 g zinc sulphate. This may be because of the increment of biomass production by zinc sulphate resulting in dilution effects and thus, phosphorus concentration gets reduced. The reason may also be the antagonistic reactions between zinc and phosphorus. The results are in confirmation with the findings of (6) who told that the concentrations of phosphorus were low in the samples from spray treatments of zinc sulphate. In contrary to it, (5) reported that soil and foliar spray combinations of N + Zn treatment in apple had relatively higher leaf phosphorus. Ferrous sulphate also significantly affected phosphorus content in leaves. Maximum content (0.17%) was observed under control treatment as well as plants receiving 10 g and 20 g ferrous sulphate, while minimum (0.16%) was noted from plants receiving 30 g ferrous sulphate. Phosphorus content in guava leaves did not vary significantly due to interaction between zinc sulphate and ferrous sulphate.

**Potassium content in leaf :** It is amply clear from the data showed in Table-2 that zinc sulphate had significant

effect on potassium content in guava leaves. Maximum potassium content (1.61%) was observed under control treatment, which was significantly higher than all other treatments, while minimum (1.40%) was observed from the plants receiving 90 g zinc sulphate. The reason may be the increase in biomass production by zinc sulphate which leads to dilution effect resulting in reduction of potassium concentration. The results are in contradiction with the results obtained by (5) who told that soil and foliar spray combinations of N + Zn treatment had relatively higher leaf potassium, whereas non-significant effect was observed by (6) in apple. Potassium content in leaves was affected significantly with the application of ferrous sulphate also. Maximum content (1.60%) was found under control treatment, while minimum (1.37%) was observed from plants receiving 10 g ferrous sulphate. The results are more or less in accordance with the findings of (9) in Satsuma mandarin that leaf potassium content was significantly higher in the non-iron treatment tree than in the iron treatment. The interaction between zinc sulphate and ferrous sulphate had non-significant effect on potassium content in leaves.

**Zinc content in leaf :** Data from Table-3 shows that zinc content in guava leaves varied significantly due to zinc sulphate treatment. Maximum zinc content (34.34 ppm) was observed with the application of 90 g zinc sulphate, which was closely followed by 60 g zinc sulphate and significantly higher than all other treatments, while minimum (29.33 ppm) was noted under control treatment. This may be because of the reason that zinc easily moved down the soil profile and was efficiently taken up and transported to above ground parts. These findings are in close agreement with the results of (10) who told that zinc concentration in desirable foliage increased proportionally as the amount of soil-banded zinc increased in pecan orchard. Similar increasing trend was noted by (11) in (12) in Sathgudi orange, (13) in apple, (14) in papaya, (6), (15), (5) in apple and (16) in mango.

Ferrous sulphate also had significant effect on zinc content in leaves. Maximum zinc content (33.63 ppm) was found with application of 20 g ferrous sulphate, which was significantly higher than all other treatments while, minimum (30.36 ppm) was noted from plants under control treatment. Zinc and iron were found to be inter-competitive in leaves and the increase of leaf iron concentration could reduce zinc concentration. Earlier such findings were reported by (9) in Satsuma mandarin that leaf zinc concentrations were significantly higher in the non-iron treatments than that in the iron treatment. The interaction between zinc sulphate and ferrous sulphate gave non-significant results for zinc concentration in leaves.

**Table-1 : Effect of zinc sulphate and ferrous sulphate on total nitrogen (%) and total phosphorus (%) content of guava leaf cv. Hisar Safeda.**

ZnSO <sub>4</sub> (g/plant)	FeSO <sub>4</sub> (g/plant)									
	Total nitrogen (%)					Total phosphorus (%)				
	0	10	20	30	Mean	0	10	20	30	Mean
Zero	1.66	1.67	1.69	1.68	1.68	0.19	0.18	0.18	0.17	0.18
30	1.71	1.73	1.76	1.74	1.73	0.17	0.17	0.17	0.16	0.17
60	1.68	1.70	1.74	1.72	1.71	0.17	0.16	0.16	0.16	0.16
90	1.72	1.73	1.76	1.74	1.74	0.17	0.17	0.16	0.17	0.17
Mean	1.69	1.71	1.74	1.72		0.17	0.17	0.17	0.16	
CD at 5%	Zn= 0.02, Fe= 0.02, Zn X Fe= NS					Zn= 0.01, Fe= 0.01, Zn X Fe= NS				

**Table-2 : Effect of zinc sulphate and ferrous sulphate on total potassium (%) content of guava leaf cv. Hisar Safeda.**

ZnSO <sub>4</sub> (g/plant)	FeSO <sub>4</sub> (g/plant)				
	Zero	10	20	30	Mean
Zero	1.77	1.49	1.55	1.62	1.61
30	1.65	1.34	1.45	1.54	1.49
60	1.55	1.33	1.42	1.44	1.44
90	1.45	1.33	1.40	1.42	1.40
Mean	1.60	1.37	1.46	1.51	
CD at 5%	Zn= 0.07, Fe= 0.07, Zn x Fe= NS				

**Table-3 : Effect of zinc sulphate and ferrous sulphate on total zinc content (ppm) and total iron content (ppm) content of guava leaf cv. Hisar Safeda.**

ZnSO <sub>4</sub> (g/plant)	FeSO <sub>4</sub> (g/plant)									
	Total zinc content (ppm)					Total iron content (ppm)				
	0	10	20	30	Mean	0	10	20	30	Mean
Zero	27.20	28.13	31.36	30.60	29.33	160.0	168.5	170.4	172.0	167.7
30	28.84	30.71	32.67	31.14	30.84	166.7	168.7	170.2	172.0	169.4
60	32.23	33.20	34.81	33.84	33.52	163.7	165.9	167.8	176.7	168.5
90	33.18	33.82	35.67	34.71	34.34	164.9	166.6	168.7	175.7	169.0
Mean	30.36	31.47	33.63	32.57		163.8	167.4	169.3	174.1	
CD at 5%	Zn= 0.87, Fe= 0.87, Zn X Fe= NS					Zn= 0.01, Fe= 0.01, Zn X Fe= NS				

**Iron content in leaf :** Persual from the data shown in Table-3 indicates that zinc sulphate had significant effect on iron content in leaves. Maximum iron content (169.4 ppm) was observed from the plants receiving 30 g zinc sulphate, whereas minimum (167.7 ppm) was noted under control treatment. This may be because the iron was immobilized somewhere in the leaf in an unavailable form. Similar increase in iron concentration of leaf with iron treatment had been observed by (9) in Satsuma mandarin that leaf iron concentrations were less in the non-iron treatments than that in the iron treatment. Iron content in leaves was significantly affected with the application of ferrous sulphate also. Maximum iron (174.1 ppm) was observed from the plants receiving 30 g ferrous sulphate, which was significantly higher than all other treatments, while minimum (163.8 ppm) was noted from control treatment. No significant effect was found on iron content in leaves due to interaction between zinc sulphate and ferrous sulphate.

## References

1. Wolf B. (1982). A comprehensive system of leaf analyses and its use for diagnosing crop nutrient status. *Communications in Soil Science and Plant Analysis*, 3(12): 1035-1059.
2. Rampal Singh, Ompal Singh and H.S. Rathore (2020). Foliar fertilization of vegetable and fruit plants. *Frontiers in Crop Improvement*, 8(1): 1-10.
3. Lindner R.C. (1944). Rapid analytical methods for some of the more common inorganic constituents of plant tissues. *Plant physiology*, 19(1): 76.
4. Koenig R. and Johnson C. (1942). Colorimetric determination of phosphorus in biological materials. *Industrial & Engineering Chemistry Analytical Edition*, 14(2): 155-156.
5. Amiri M.E., Fallahi E. and Golchin A. (2008). Influence of foliar and ground fertilization on yield, fruit quality, and soil, leaf, and fruit mineral nutrients in apple. *Journal of plant nutrition*, 31(3): 515-525.
6. Rasouli Sadaghiani M.H., Malakouti M.J. and Samar S.M.

- (2002). The effectiveness of different application methods of zinc sulfate on nutritional conditions of apple in calcareous soils of Iran.
7. Shewy A.A. and Abdel-Khalek A.I. (2014). Physiological studies on the effect of foliar sprays with some micronutrients on leaf mineral content, yield and fruit quality of "Florida Prince and Desert Red" peach trees. *Trends in Horticultural Research*, 4(1): 20-30.
  8. Domenico Rombolà A., Toselli M., Carpintero J., Ammari T., Quartieri M., Torrent J. and Marangoni B. (2003). Prevention of iron-deficiency induced chlorosis in kiwifruit (*Actinidia deliciosa*) through soil application of synthetic vivianite in a calcareous soil. *Journal of plant nutrition*, 26(10-11): 2031-2041.
  9. Huang H., Hu C.X., Tan Q., Hu X., Sun X. and Bi L. (2012). Effects of Fe-EDDHA application on iron chlorosis of citrus trees and comparison of evaluations on nutrient balance with three approaches. *Scientia Horticulturae*, 146: 137-142.
  10. Wood B.W. (2007). Correction of zinc deficiency in pecan by soil banding. *Hort. Science*, 42(7): 1554-1558.
  11. Nijjar G.S. and Brar S.S. (1977). Comparison of Soil and Foliar Applied Zinc in Kinnow. *Indian Journal of Horticulture*, 34(2): 130-136.
  12. Devi D.D., Srinivasan P.S. and Balkrishnan K. (1997). Influence of Zn, Fe and Mn on photosynthesis and yield of *Citrus sinensis*. *Indian Journal of Plant Physiology*, 2(2): 174-176.
  13. Pavan M.A. (1998). Response of apple to soil applied zinc. *Pesquisa Agropecuaria Brasileira* (Brazil).
  14. Kavitha M., Kumar N., Jeyakumar P. and Soorianathasundaram K. (2002). Changes in the nutrient status of papaya cv. CO-5 as influenced by zinc and boron application. *South Indian Horticulture*, 50(1/6): 200-206.
  15. Wojcik P. (2007). Vegetative and reproductive responses of apple trees to zinc fertilization under conditions of acid coarse-textured soil. *Journal of plant nutrition*, 30(11): 1791-1802.
  16. Bahadur L., Malhi C.S. and Singh Z. (1998). Effect of foliar and soil applications of zinc sulphate on zinc uptake, tree size, yield, and fruit quality of mango. *Journal of plant nutrition*, 21(3): 589-600.